

Delphi Technique Theoretical Framework in Qualitative Research

¹, Arash Habibi, ², Azam Sarafrazi, ³, Sedigheh Izadyar

¹ Master of Business Administration, Isfahan University, Iran

² Master of EMBA, Payam Noor University, Iran

³ Master of Marketing Management, Guilan University, Iran

ABSTRACT

Using specialized knowledge and perspectives of a set in decision-makings about issues that are qualitative is very helpful. Delphi technique is a group knowledge acquisition method, which is also used for qualitative issue decision-makings. Delphi technique can be used for qualitative research that is exploratory and identifying the nature and fundamental elements of a phenomenon is a basis for study. It is a structured process for collecting data during the successive rounds and group consensus. Despite over a half century of using Delphi in scientific and academic studies, there are still several ambiguities about it. The main problem in using the Delphi technique is lack of a clear theoretical framework for using this technique. Therefore, this study aimed to present a comprehensive theoretical framework for the application of Delphi technique in qualitative research. In this theoretical framework, the application and consensus principles of Delphi technique in qualitative research were clearly explained.

Keywords - Delphi Technique, Qualitative Research, Theoretical Framework

Date of Submission: 07 April 2014

Date of Publication: 25 April 2014

I. INTRODUCTION

Delphi is a name of temple in ancient Greek, which is the origin of Apollo's divine inspirations, god of sun and music. Greeks and ancient nations came in Pythia, the goddess of Delphi temple to predict the future [1] [2]. The Delphi technique was originally proposed based on people's conjecture, judgment, and inspiration but gradually took the academic form. For the first time in the late 1950s, in a research by U.S. RAND Corporation, the Delphi was introduced for the scientific study of experts' opinions on military defense project. However, for the security reasons, this technique was not proposed over ten years and in 1963, Dalkey and Helmer introduced it [3] [4]. Its first non-military use was suggested for economic development planning [5] [6] [7]. This technique gradually found its place in the academic studies but from the mid-90s, it became highly popular. Landeta's study indicated that since 1995 to 1999 a total 444 articles on this technique have been published in "Science Direct" and "ABI / Inform" journals. Since 2000 to 2004, this number has increased to 667 articles [6]. Among the various features of the Delphi technique, its four features are usually unchanged including anonymity, iteration, controlled feedback, and statistical "group response" [8] [9] [10].

Most researchers have used all [11] or some [12] [13] of the definition set out by Linstone and Turoff (1975) who define the Delphi technique as " a method for structuring a group communication process so that the process is effective in allowing a group of individuals, as a whole, to deal with a complex problem" (p. 3) [14]. The main purpose of the Delphi method is 'to acquire the most reliable consensus of a group of experts' opinion by 'a series of intensive questionnaires combined with controlled opinion feedback' (p. 458) [15]. By obtaining the consensus of a group of experts using the process, researchers can identify and prioritize issues and develop a framework to recognize them [16] [2]. Hasson et al. (2000) argued that modified Delphi, policy Delphi, and real-time Delphi have all been used, while there are many potential types of Delphi techniques [17]. Three broad categories are generally in use and these are classical, policy and decision [18]. The Delphi technique is applied as a tools and method for consensus building using a series of questionnaires for data collection from a panel of selected participants [15] [19] [14] [20] [21] [22].

Undoubtedly, the collective wisdom in decision-making can lead to more complete and comprehensive decision-making. However, the group problem solving through traditional consensus method is associated with numerous difficulties. The group members who have high self-confidence are dominating the weaker members. Some are also pressured to conform to group comments and or because they respect other people, they do not express their opinions. Hence, a group problem solving approach will be often futile and ineffective. The anonymity principle is used to solve this problem. In Delphi technique, the experts and people who are used the survey do not know each other. Anonymity ensures overcoming the obstacles of groupthink [23] [24] [25] [1] [26]. A coordinator collects the experts' opinions and then he/she provides other members with the summarized results. Then, based on the summarized results in the previous step, individuals again adjust and express their opinions. Finally, after reaching a consensus, the results are discussed in terms of a statistical report (usually mean or median) and are used for decision-making [9] [27] [28].

The Delphi technique is a research approach to gain consensus using a series of questionnaires and the provision of feedback to participants who have expertise in key areas. The main problem in using the Delphi technique is the lack of a clear theoretical framework for using this technique. Therefore, this study aimed to present a comprehensive theoretical framework for the application of Delphi technique in qualitative research. In this theoretical framework, we clearly explain the application and consensus principles of Delphi technique in qualitative research.

II. DELPHI TECHNIQUE THEORETICAL FRAMEWORK

The major weakness of Delphi is the lack of a theoretical framework. Delphi as a research methodology has been variously presented as a survey, study, procedure, method, approach, polling, and technique [14] [29] [30] [31] [24] [25] [32] [33] [11] [34] [35] [36] [37] [38]. Nevertheless, it is generally acknowledged that the Delphi methodology is not uniform in its application [17] [1]. In addition, there are always ambiguities about the application requirements, panel size, how to select the panel, and recognizing the end of Delphi stages. In this comparative study, a framework is presented for the application of the Delphi technique in the qualitative decision-makings.




Fig 1. Theoretical framework of Delphi technique in qualitative research

2.1 Delphi Technique's Application Requirements

The most important requirements for the use of Delphi are the need for experts' judgment, group consensus to achieve the results, anonymity in data collection, a complex, multidimensional, and interdisciplinary problem, lack of consensus and imperfect knowledge, experienced and capable experts, dispersion of experts, no time limitation, and lack of cost-effective method [39] [6] [40] [28] [7] .

In qualitative research that is mainly based on the individuals' judgments and opinions, there is a prerequisite for using the Delphi method. If these studies are based on the expert opinions, then the use of inferential statistical techniques such as mean tests will not be justified. Since either experts are not so much that can be accessed wide samples of them or the time and cost to access them are difficult. In addition, to review the experts' opinions this proverb is truer: "you may know by a handful the whole sack".

Therefore, the two types of qualitative research should be distinguished to use the Delphi technique. Delphi technique can be used for qualitative research that is exploratory and identifying the nature and fundamental elements of a phenomenon is a basis for the study. Consider the problem of measuring customer satisfaction. If you were to examine the customer satisfaction, statistical sampling and inferential statistical methods would be used. However, if you decide to measure the customer satisfaction, then experts' opinions and the Delphi technique can be used. One of main application of Delphi technique is screening the items in operations research problems and of multi-criteria decision-making (MCDM) techniques.

2.2 Composition and Panel Size

Essentially, an accurate mechanism does not exist for identifying the number of individuals or the number of panels for inclusion in any individual study [41]. Although there are some disagreements about the composition and panel size of Delphi technique, a dominant pattern can be detected. It has been recommended that the panel size may vary according to the topics covered, the nature of different viewpoints included, and the time and money available [18] and it is also suggested to use a combination of individuals with multiple specialties and heterogeneous groups better than the homogeneous groups [1] [26]. Hsu and Sandford (2007) claimed that, "Delphi subjects should be highly trained and competent within the specialized area of knowledge related to the target issue" (p.4) [42]. Hogarth (1978) argued that between six and twelve members are ideal for Delphi technique and according to Clayton (1997), if a mixture of experts with different specialties is used, between five and ten members are sufficient (p. 5) [26]. While some Delphi studies considered fewer than 10 members in their panels [43] [44], other studies included more than 100 participants [45] [46].

Delphi panel is done with the participation of individuals who have the knowledge and expertise of the study subject. These people are known as the Delphi panel. One of the most important phases of Delphi technique is selecting eligible members for the Delphi panel because the validity of the results depends on the competence and knowledge of panel members [1]. In previous studies, Delphi has not recommended a specific method for sampling. Snowball technique can be used for sampling. The snowball sampling is one of the most common approaches of sequential sampling. This type of sampling is a non-probability method, which is also randomly selected. This method is suitable when the members of a group or community cannot be easily identified. In this method, the researcher first identifies some people and after receiving the information, he/she wants them to introduce other people [47]. This method can be also used to identify the experts in a particular field [48].

2.3 Gathering Expert Opinions

In qualitative research, which aims to determine the importance or screening of items, Likert scale can be used to gather the experts' opinions. Five-point or seven-point Likert scale is common. To develop five-point Likert scale, two "extremely agree" and "extremely disagree" linguistic scales are used at both ends of a spectrum. By defining the intermediate values, this scale can be developed and nine-point scales can be also used.

Diefenbach, Weinstein, and O'Reilly (1993) conducted an investigation of a range of Likert items, including 2-point, 5-point, 7-point, 9-point, 11-point, 12-point, and percentage (100-point) varieties [49]. Lewis (1993) found that 7-point scales resulted in stronger correlations with t-test results [50]. It was demonstrated that "no scale performed significantly better than the seven-point verbal category scale on any criterion" in the two studies carried out (p. 189) [49].



Fig 2. A five-point Likert scale [51]




Fig 3. A seven-point Likert scale [51]

After gathering the experts' opinions, the mean score of their opinions on each dimension is calculated. Given the theoretical framework, if there is no consensus, the experts will be provided with the calculated mean as a controlled feedback with questionnaire. After several rounds, when the consensus was achieved, based on the average of the final round, the items are screened. If nine-point scale is used, typically, the criteria with mean less than seven are removed. For seven-point and five-point scales, criteria with means lower than five and four are respectively removed.

2.4 Determining the Level of Consensus

Another problem with the Delphi technique is a scientific method for determining the level of consensus. A variety of methods has been proposed in different studies. Some researchers have determined Delphi rounds as a basis for reaching agreement. Between 2 and 10 rounds have been reported in various articles [53]. Fan and Cheng have believed that research has shown that three rounds are adequate for the Delphi technique [40]. Some also argued that the traditional Delphi contains four rounds that researchers have shortened it into two or three rounds to achieve the study's objectives [54] [5]. Such an approach is a target displacement. In fact, reaching a consensus is a basis for the end of Delphi rounds while in this approach doing a certain number of rounds is regarded as a basis for reaching an agreement. So, another solution must be identified to reach consensus. In a study, according to 114 articles about the Delphi technique, von der Gracht (2012) has presented 15 ways to achieve consensus [9].

Kendall's coefficient of concordance (Kendall's W), a consensus criterion representing the level of consensus between the participants [55] [56] [57], was calculated along with the mean rank and standard deviation. Kendall's coefficient of concordance ranges from 0 to 1, indicating the degree of consensus reached by the panel (strong consensus for $W > 0.7$; moderate consensus for $W = 0.5$; and weak consensus for $W < 0.3$) [58]. Kendall's coefficient of concordance is also a scale for determining the level of coordination and agreement between several ranks of n phenomenon. This scale shows a rank correlation between m ranking set. It can be used to determine the inter-judge reliability. Kendall's coefficient of concordance indicates that those who have ordered several categories according to their importance have used the same criteria to judge the importance of each category and in this regard, they agree to each other [57]. Kendall's coefficient is calculated as follows:

$$W = \frac{12S}{m^2(n^2 - n)} \tag{1}$$

Where

$$S = \sum_{i=1}^n (R_i - \bar{R})^2$$

- R_i = total ranks of a factor
- m = number of rank sets or judges
- n = number of ranked factors or phenomena

The value of this scale during the full coordination or agreement is one and in the absence of perfect coordination, it is zero. Schmidt (1997) proposed two statistical criteria to make decisions about stopping or continuing the Delphi rounds. The first criteria is a strong consensus among the panel members which is determined based on Kendall's coefficient of concordance. If there is no such a consensus, the constant coefficient or its negligible growth in two successive rounds shows that consensus has not been increased and the survey process must be stopped. Statistically significant coefficient w is not enough to stop a Delphi process. For panels with fewer than ten members, very small values of w are also significant [56].

III. CONCLUSION

To summarize, the Delphi technique is a research approach to gain consensus using a series of questionnaires and the provision of feedback to participants who have expertise in key areas. This method is especially useful when researchers need to collect ideas from isolated experts on a specific topic and establish agreement to discover the underlying assumptions or perspectives among the experts. However, the major weakness of Delphi is the lack of a theoretical framework. Delphi as a research methodology has been variously presented as a survey, study, procedure, method, approach, polling, and technique. Nevertheless, it is generally acknowledged that the Delphi methodology is not uniform in its application.

Although previous studies have always been disadvantaged by the lack of a theoretical framework for Delphi technique, considering numerous studies, access to a theoretical framework is simple. Given the findings of this study, it can be said that in qualitative research, it is recommended to use a group of ten experts with different specialties. The snowball sampling technique can also be used to identify and select the sample. If the purpose is to summarize and screen an item set, a nine-point Likert scale can be used to gather the experts' opinions. Kendall's coefficient of concordance can be utilized to determine the agreement and end of the Delphi technique. Therefore, it can be concluded that the Delphi technique is an appropriate and acceptable mechanism to achieve consensus when developing sets of indicators.

REFERENCES

- [1] Powell, C. (2003). The Delphi technique: Myths and realities. *Journal of Advanced Nursing*, 41(4), 376-382.
- [2] Okoli C and SD Pawlowski (2004) The Delphi method as a research tool: An example, design considerations, and applications. *Information & Management* 42(1), 15–29.
- [3] Kauko, Karlo., Palmroos, Peter. (2014) The Delphi method in forecasting financial markets An experimental study. *International Journal of Forecasting* 30:2, 313-327 Online publication date: 1-Apr-2014.
- [4] Keil, Mark. Hyung Koo Lee, Tianjie Deng. (2013) Understanding the most critical skills for managing IT projects: A Delphi study of IT project managers, *Information & Management*, Volume 50, Issue 7, November 2013, Pages 398-414
- [5] Turoff M, Linstone HA. The Delphi method: techniques and applications. [Cited 2008 Oct 19]. Available from: <http://is.njit.edu/pubs/delphibook/>
- [6] Landeta J., (2006), "Current validity of the Delphi method in social sciences", *Technological Forecasting and Social Change*, Vol. 73, No. 5, pp: 467-82.
- [7] Meijering J.V., J.K. Kampen, H. Tobi (2013), Quantifying the development of agreement among experts in Delphi studies ,*Technological Forecasting and Social Change*, Volume 80, Issue 8, October 2013, Pages 1607-1614
- [8] Rowe, G., & Wright, G., Expert opinions in forecasting: the role of the Delphi technique, in: J.S. Armstrong (Ed.), *Principles of Forecasting: A Handbook for Researchers and Practitioners*, Kluwer Academic Publishers, Boston et al., 2001, pp. 125–144.
- [9] von der Gracht, Heiko, Consensus measurement in Delphi studies Re-opinion and implications for future quality assurance, *Technological Forecasting & Social Change* 79 (2012) 1525–1536.
- [10] Jünger, Saskia., Sarah Brearley, Sheila Payne, Aukje K. Mantel-Teeuwisse, Thomas Lynch, Willem Scholten, Lukas Radbruch (2013), Consensus Building on Access to Controlled Medicines: A Four-Stage Delphi Consensus Procedure , *Journal of Pain and Symptom Management*, Volume 46, Issue 6, December 2013, Pages 897-910
- [11] Wang, C.C., Wang, Y., Zhang, K., Fang, J., Liu, W., Luo, S., Tang, S., Wang, S. and Lu, V.C. (2003) Reproductive Health Indicators for China's Rural Areas. *Social Science and Medicine*, 57(2): 217-225.
- [12] Gupta, U.G. and Clarke, R.E. (1996) Theory and Applications of the Delphi Technique: A Bibliography (1975-1994) *Technological Forecasting and Social Change*, 53: 185-211
- [13] Robertson, H.A. and MacKinnon, N.J. (2002) Development of a List of Consensus-Approved Clinical Indicators of Preventable Drug-Related Morbidity in Older Adults. *Clinical Therapeutics*, 24(10): 1595-1613.
- [14] Linstone, H.A. and Turoff, M. (eds.) (1975) *the Delphi Method Techniques and Applications*, Massachusetts, Reading: Addison-Wesley.
- [15] Dalkey N and O Helmer (1963) An experimental application of the Delphi method to the use of experts. *Management Science* 9(3), 458–467
- [16] Greatorex J and T Dexter (2000) An accessible analytical approach for investigating what happens between the rounds of a Delphi study. *Journal of Advanced Nursing* 32(4), 1016–1024.
- [17] Hasson, F., Keeney, S. and McKenna, H. (2000) Research Guidelines for the Delphi Survey Technique. *Journal of Advanced Nursing*, 32(4): 1008-1015.
- [18] van Zolingen, S.J. and Klaassen, C.A. (2003) Selection Processes in a Delphi Study about Key Qualifications in Senior Secondary Vocational Education. *Technological Forecasting and Social Change*, 70: 317-340.
- [19] Dalkey, N. C. (1969). An experimental study of group opinion. *Futures*, 1 (5), 408-426.
- [20] Lindeman, C. A. (1981). Priorities within the health care system: A Delphi survey. Kansas City, MO: American Nurses' Association.
- [21] Martino, J. P. (1983). *Technological forecasting for decision making*. New York: North-Holland.

- [22] Young, S. J., & Jamieson, L. M. (2001). Delivery methodology of the Delphi: A comparison of two approaches. *Journal of Park and Recreation Administration*, 19 (1), 42-58.
- [23] Mitroff, I. I., & Turoff, M. (1975). Philosophical and methodological foundations of Delphi. In H. A. Linstone & M. Turoff (Eds.), *The Delphi method: Techniques and applications* (pp. 17-34). Boston, MA: Addison-Wesley.
- [24] Browne N, Robinson L, Richardson A. A Delphi study on the research priorities of European oncology nurses. *Eur J Oncol Nurs* 2002 Sep; 6(3): 133-44.
- [25] McKenna H, Hasson F, Smith M. A Delphi survey of midwives and midwifery students to identify non-midwifery duties. *Midwifery* 2002 Dec; 18(4): 314-22.
- [26] Somerville, Jerry A (2008). *Effective Use of the Delphi Process in Research: Its Characteristics, Strengths, and Limitations*
- [27] Antcliff D, Keeley P, et al. (2013). The development of an activity-pacing questionnaire for chronic pain and/or fatigue: a Delphi technique, *Physiotherapy*
- [28] Cowan, D., et al. Direct care activities for assistants in nursing in inpatient mental health settings in Australia : A modified Delphi study. *Collegian* (2013), <http://dx.doi.org/10.1016/j.colegn.2013.11.003>
- [29] Crisp, J., Pelletier, D., Duffield, C., Adams, A. and Nagy, S. (1997) *The Delphi Method?* *Nursing Research*, 46(2): 116-118
- [30] Broomfield, D. and Humphries, G.M. (2001) Using the Delphi Technique to Identify the Cancer Education Requirements of General Practitioners. *Medical Education*, 35: 928-937.
- [31] Sharkey, S.B. and Sharples, A.Y. (2001) An Approach to Consensus Building Using the Delphi Technique: Developing a Learning Resource in Mental Health. *Nurse Education Today*, 21: 398-408.
- [32] Rogers, M.R. and Lopez, E.C. (20 02) Identifying Critical Cross-Cultural School Psycholog Competencies. *Journal of School Psychology*, 40(2): 115-141
- [33] Snyder-Halpern, R. (2002) Indicators of Organizational Readiness for Clinical Information Technology Systems Innovation: A Delphi Study. *International Journal of Medical Informatics*, 63: 179-204.
- [34] Campbell, S.M., Shield, T., Rogers, A. and Gask, L. (2004) How do Stakeholder Groups Vary in a Delphi Technique About Primary Mental Health Care and What Factors Influence their Ratings? *Qual Saf Health Care*, 13 (6): 428-34.
- [35] Iavicoli, S., Rondinone, B., Marinaccio, A. and Fingerhut, M. (2005) Identification of Research Priorities in Occupational Health. *Occup Environ Med*, 62 (2):71-2.
- [36] Miró J, Nieto R, Huguet A. Predictive factors of chronic pain and disability in whiplash: a Delphi poll. *Eur J Pain* 2008 Jan; 12(1): 30-47. Epub 2007 Apr 2.
- [37] Skulmosky G, Hartman FT, Krahn J. The Delphi method for graduate research. *Journal of Information Technology Education* 2007; 6: 1-21.
- [38] Langlands RL, Jorm AF, Kelly CM, Kitchener BA. First aid for depression: a Delphi consensus study with consumers, carers and clinicians. *J Affect Disord* 2008 Jan; 105(1-3): 157-65. Epub 2007 Jun 14.
- [39] Walker, AM and Selfe, James (1996) The Delphi method: a useful tool for the allied health researcher. *International Journal of Therapy and Rehabilitation*, 3 (12).
- [40] Fan, C.K, Cheng C.-L., A study to identify the training needs of life insurance sales representatives in Taiwan using the Delphi approach. *Int. J. Train. Dev.* 10 (2006) 212–226.
- [41] Williams, P.L. and Webb, C. (1994) The Delphi Technique: A Methodological Discussion. *Journal of Advanced Nursing*, 19: 180-186
- [42] Hsu, C., & Sandford, B. (2007). The Delphi technique: Making sense of consensus. *Practical Assessment, Research and Evaluation*, 12, 1–8.
- [43] Malone D, J Abarca, P Hansten, A Grizzle, E Armstrong, RV Bergen, B Duncan-Edgar, S Solomon and R Lipton (2005) Identification of serious drug-drug interactions: Results of the partnership to prevent drug-drug interactions. *American Journal of Geriatric Pharmacotherapy* 3(2), 65–76.
- [44] Strasser S, L London and E Kortenbout (2005) Developing a competence framework and evaluation tool for primary care nursing in South Africa. *Education for Health* 18(2), 133–144.
- [45] Kelly KP and D Porock (2005) A survey of pediatric oncology nurses' perceptions of parent educational needs. *Journal of Pediatric Oncology Nursing* 22(1), 58–66
- [46] Meadows A, LMaine, E Keyes, K Pearson and K Finstuen (2005) Pharmacy executive leadership issues and associated skills, knowledge, and abilities. *Journal of the American Pharmacists Association* 45(1), 55–62.
- [47] Babbie E. *The basics of social research*. 2nd ed. ed. Belmont, Calif. ; London: Wadsworth/Thomson Learning; 2002.
- [48] Macnee CL, McCabe SRN. *Understanding nursing research: using research in evidence-based practice*. 2nd ed. ed. Philadelphia, PA: Wolters Kluwer Health/Lippincott Williams & Wilkins; 2008
- [49] Diefenbach, M.A., Weinstein, N.D., & O'Reilly, J. (1993). Scales for assessing perceptions of health hazard susceptibility, *Health Education Research*, 8, 181-192.
- [50] Lewis, J. R. (1993). Multipoint scales: Mean and median differences and observed significance levels. *International Journal of Human-Computer Interaction*, 5(4), 383-392.
- [51] Likert, R. (1932). A technique for the measurement of attitudes. *Archives of Psychology*, 22(140), 1–55.
- [52] Fry M, Burr G. Using the Delphi technique to design a self-reporting triage survey tool. *Accid Emerg Nurs* 2001 Oct; 9(4): 235-41.
- [53] Windle PE. Delphi technique: assessing component needs. *J Perianesth Nurs* 2004 Feb; 19(1): 46-7.
- [54] Nevo D and YE Chan (2007) A Delphi study of knowledge management systems: Scope and requirements. *Information & Management* 44(6), 583–597.
- [55] Schmidt R.C.; *Managing Delphi surveys using nonparametric statistical techniques; Decision Sciences*, Vol.28, No.3, 1997.
- [56] Siegel, S., and Castellan, N. J., Jr., 1988, "Nonparametric statistics for the behavioral sciences (2nd ed.)", New York: McGraw-Hill.
- [57] Schmidt RC, K Lyytinen, M Keil and P Cule (2001) Identifying software project risks: An international Delphi study. *Journal of Management Information Systems* 17(4), 5–36.